Syntopy in California red-legged and foothill yellow-legged frogs in their aquatic habitat

December 29, 2023

RESEARCH NOTE

Jeff A. Alvarez[1]*, Mikhela Aiken[2], and Jeffery T. Wilcox[3]

[1] *The Wildlife Project, P.O. Box 188888, Sacramento, CA 95818, USA*
https://orcid.org/0000-0003-1875-6327
[2] *Area West Environmental, 6248 Main Avenue, Suite #C, Orangevale, CA 95662, USA*
[3] *Mitsui Ranch Preserve, 3124 Sonoma Mountain Road, Petaluma, CA 95954, USA*
https://orcid.org/0000-0002-3271-8514

*Corresponding Author: *jeff@thewildlifeproject.com*

Published 29 December 2023 • doi.org/10.51492/cfwj.109.16

Key words: aquatic, California red-legged frog, declining, foothill yellow-legged frog, management, protected, *Rana boylii*, *Rana draytonii*, range, sympatry, threatened

Citation: Alvarez, J. A., M. Aiken, and J. T. Wilcox. 2023. Syntopy in California red-legged and foothill yellow-legged frogs in their aquatic habitat. California Fish and Wildlife Journal 109:e16.

Editor: Jennifer Olson, Watershed Restoration Grants Branch

Submitted: 13 July 2023; **Accepted:** 5 September 2023

Copyright: ©2023, Alvarez et al. This is an open access article and is considered public domain. Users have the right to read, download, copy, distribute, print, search, or link to the full texts of articles in this journal, crawl them for indexing, pass them as data to software, or use them for any other lawful purpose, provided the authors and the California Department of Fish and Wildlife are acknowledged.

Competing Interests: The authors have not declared any competing interests.

Co-occurrence of two or more species of amphibians with biphasic life cycles is extremely common in North America, particularly around the east coast of the United States (Petranka 1998; Lannoo 2005; McGinnis and Stebbins 2018). On the west coast, different amphibians are often sympatric (share the same distribution) and/or syntopic (share the same habitat at the same time) over a wide range of genera and species, including various assemblages of *Ambystoma, Taricha, Dicamptodon*, *Pseudacris, Spea, Anaxyrus, Rana, Lithobates,* and others (Storer 1925; Petranka 1998; Lannoo 2005; McGinnis and

Stebbins 2018; Flaxington 2021). For example, the California giant salamander (*Dicamptodon ensatus*) and northwestern salamander (*Ambystoma gracile*) are commonly found under the same cover objects and often breed in the same aquatic habitats (Storer 1925; Stebbins 2003). The wide-ranging rough-skinned newt (*Taricha granulosa*) and California newt (*T. torosa*) are frequently found together in both terrestrial and aquatic breeding habitat where the species are syntopic in the southern portion of the range of *T. granulosa* (Stebbins 2003; Stebbins and McGinnis 2012). Similarly, in Santa Cruz County, the California tiger salamander (*Ambystoma californiense*) and its congener the Santa Cruz long-toed salamander (*A. macrodactylum croceum*) are syntopic (J. Alvarez, pers. obs.). Recently, Alvarez et al. (2013) reported a wide area where the California tiger salamander and California red-legged frog (*Rana draytonii*) were synoptic.

California red-legged frogs are listed as threatened under the federal Endangered Species Act and are an Amphibian Species of Special Concern in California (Thomson et al. 2016). The California red-legged frog is closely associated with aquatic freshwater habitats surrounded by grasslands, chaparral, woodlands, and forest habitat types (Storer 1925; Hayes and Jennings 1988; Bulger et al. 2003; Allaback et al. 2010). The species is frequently syntopic with other native amphibians in their aquatic breeding habitat, including California newt, rough-skinned newt, California tiger salamander, Pacific treefrog (*Hyliola regilla*), western toad (*Anaxyrus boreas*), as well as non-native American bullfrog (*Lithobates catesbeianus*; Storer 1925; Feaver 1971; Hayes and Tennant 1985; Rathbun 1998; Cook and Jennings 2007; Alvarez et al. 2013).

Foothill yellow-legged frogs (*R. boylii*) were a California Amphibian Species of Special Concern since 1994 (Jennings and Hayes 1994), were recently considered candidate species for federal listing (USFWS 2022), and several clades were listed in 2020 under the California Endangered Species Act as threatened or endangered (CFGC 2020). The foothill yellow-legged frog is another Pacific Coast ranid species but is better known for being associated with both perennial and intermittent creeks, rivers, and streams (Zwiefel 1955; Bourque 2008; Bondi et al. 2013; McGinnis and Stebbins 2018). Zwiefel (1955) noted that he rarely found foothill yellow-legged frogs more than a single hop from a creek or stream. More recently Wilcox and Alvarez (2019) and Alvarez and Wilcox (2021a) reported the foothill yellow-legged frogs occasionally use lentic systems for breeding, refuge, and presumably foraging.

The range of California red-legged frogs and foothill yellow legged frogs appear to have intermittent overlapping areas ranging from Mendocino and Plumas counties in the north and extending south into the Sierra Nevada and along coastal California to Baja California in Mexico (**Fig. 1**; Thomson et al. 2016; McGinnis and Stebbins 2018; Flaxington 2021). When declining species are syntopic and also protected by state or federal law, as in the case of these to native ranids, management implications may be considerable. This is particularly true when management activities require actions within aquatic breeding sites for one of the listed species and not the other. If management activities are focused on a single species without regard to the other, the actions may result in changes to habitat suitability or population persistence for one or more life stages, or direct or indirect harm of the untargeted syntopic species (Alvarez et al. 2013). Although the general belief is that the California red-legged frog and foothill yellow-legged frogs would rarely overlap due to niche differentiation (Thomson et al. 2016), we note several locations where the species co-occur and breed, seek refuge, and actively forage in the same location at the same time (i.e., syntopy). We report here numerous accounts of syntopy in the breeding habitat of two protected amphibians that have not been reported elsewhere, which may affect management of both species.




Figure 1. The ranges of California red-legged frog (CRLF), and foothill yellow-legged frog (FYLF),

areas of range overlap, and the regions of our sites of investigation in California, 2023. Geographical distributions of CRLF and FYLF were adapted from Stebbins 2003, McGinnis and Stebbins 2018, and Flaxington 2021.

We investigated the occurrence of syntopy in these two species within their ranges in California via both data we collected during our own amphibian surveys and data compiled from recent and historic records. Our observational data were collected during various independent amphibian survey projects over a large area, from spring 2000 to fall 2023 (Fig. 1; Table 1). Observational data were typically collected during other long-term monitoring efforts that included predator control, amphibian population monitoring, presence/lack of presence surveys, and other survey efforts that focused on aquatic habitat for ranids. Twenty-nine percent of the data were generated from sites where we worked two or more years and where we visited sample sites multiple times. Our surveys were timed such that both species were known to be (or presumed to be, based on previous knowledge of the sites or data from comparable sites) in a detectable life stage. This would also include when neither species was expected to be occupying upland habitats. We typically conducted surveys March through July, when breeding behavior and larval specimens were likely to be associated with aquatic sites, which occurred prior to drying of most ephemeral sites.

Table 1. Location of observed sympatry of foothill yellow-legged frog (*Rana boylii*) and California red-legged frog (*R. draytonii*) in California. Numbers following California Natural Diversity Database (CNDDB) in Source column are occurrence numbers assigned to particular observations.

Type of Observation	County	Creek/ Watershed	Aquatic Habitat Type[a]	Years syntopic[b]	Year(s) reported/ observed	Source
Current	Alameda	Arroyo Mocho	Ephemeral	3	1999, 2002, 2016	CNDDB (474, 824, 789
Current	Alameda	Upper Alameda Creek	Ephemeral	2	1997, 2014	pers. obs.
Current	Napa	Wragg Creek	Ephemeral	4	2020–2023	pers. obs.
Current	San Joaquin	Corral Hollow Creek	Ephemeral	2	1998, 2014	pers. obs
Current	Santa Clara	Arroyo Hondo	Perennial	5	2005, 2009, 2013–2015	CNDDB (1336, 1492, 196, 1490)
Current	Santa Clara	Coyote Creek	Ephemeral	1	1986, 2017	pers. obs.; Gonslin 2010
Current	Santa Clara	Llagas Creek	Perennial	1	2016	CNDDB (169, 1420)

Type of Observation	County	Creek/ Watershed	Aquatic Habitat Type[a]	Years syntopic[b]	Year(s) reported/ observed	Source
Current	Santa Clara	Upper Penitecia Creek	Ephemeral	Inf	2010	CNDDB (1337, 410)
Current	Sonoma	Carriger Creek	Perennial	1	2002	pers. obs.
Current	Sonoma	Copeland Creek	Ephemeral	12	2002, 2011–2023	pers. obs.; Alvarez and Wilcox 2021a
Current	Sonoma	Turtle Pond[c]	Perennial pond	1	2020	pers. obs.; Alvarez and Wilcox 2021a
Current	Sonoma	Nolan Creek	Perennial	1	2011	CNDDB (1338, 1474)
Current	Sonoma	Mason Creek	Ephemeral	1	2022	pers. obs.
Current	Sonoma	Sheephouse Creek	Perennial	2	2012, 2017	CNDDB (1828, 1093)
Historic[d]	El Dorado	North Fork Cosumnes River	Perennial	inf	1942	CNDDB (1923, 1377)
Historic	Lake	Putah Creek	Ephemeral	1	1945	CNDDB (1704, 1706)
Historic	Merced	Los Banos Creek	Ephemeral	1	1985	CNDDB (45, 901)
Historic	Monterey	Big Creek	Perennial	1	1975	CNDDB (5239, 793)
Historic	Santa Barbara	Gaviota Creek	Perennial	inf	1940	CNDDB (2419, 388)
Historic	Santa Barbara	Mono Creek	Ephemeral	1	1940	CNDDB (2418, 1509)
Historic	Santa Barbara	Refugio Creek	Ephemeral	inf	1976	CNDDB (819, 938)

Type of Observation	County	Creek/ Watershed	Aquatic Habitat Type[a]	Years syntopic[b]	Year(s) reported/ observed	Source
Historic	Santa Clara	East Fork Coyote Creek	Ephemeral	1	2004	CNDDB (420, 798)
Historic	Santa Clara	Guadalupe Creek	Ephemeral	1	2000	CNDDB (736, 420)
Historic	Santa Clara	Grizzly Gulch	Perennial	1	2004	CNDDB (419, 792)
Historic	Santa Clara	Middle Fork Coyote Creek	Ephemeral	inf	1986	CNDDB (199, 1545)
Historic	Santa Clara	Smith Creek	Perennial	2	1950, 1975	CNDDB (56, 1547)
Historic	Santa Clara	Soquel Creek	Perennial	1	2004	CNDDB (79, 788)
Historic	Santa Clara	Stanford University	Perennial	1	1896	CNDDB (2083, 1551)
Historic	Tuolumne	Woods Creek	Perennial	1	1950	CNDDB (1955, 571)
Historic	Baja California	Woods Creek	Perennial	1	1961	Loomis 1965

[a] Habitat type at the location of observation
[b] Years syntopic is the numbers of reported or observed years species were found together; “inf” is inferred syntopy due to being present at the same site within three years of reported observations.
[c] Artificial water body
[d] Observation categorized as historic (i.e., prior to the year 2009)

We sampled both lentic and lotic habitats with a wide range of water bodies that included physical characteristics such as: perennial and ephemeral; natural and constructed; turbid to clear waters, small and large (0.004–9.8 ha); deep and shallow (0.2–10 m); low to high elevation sites (10–2070 m); sites nested within grasslands, woodlands, and chaparral; water bodies that were contained within grazed and ungrazed lands, and those supporting a range of emergent vegetation present (0–75% cover), as well as other characteristics. Our reported observations included all aquatic sites within our access area, without regard to a subjective standard for suitability. In most cases, investigators used seines and hand-held dip nets to capture larvae; occasionally post-metamorphic and adult individuals were detected visually during site visits.

In addition to our own personal observations, we conducted a search using the California Natural

Diversity Database (CNDDB) of sites that were reported as occupied by these species. Our criteria required that both species be present within a three-year period. Although both species appear to be long-lived (i.e., 15 years for both California red-legged frog [Peralta-Garcia et al. 2022] and foothill yellow-legged frog [J. Drennan, Wildlife Biologist, Kleinfelder, pers. comm.]), we used 3 years since that appeared to be the approximate average of all reported ages for both species to reach breeding age (capable of being detected at any life stage). We then categorized sites with observations older than 15 years (the presumed approximate lifespan of either species) as historic (n = 16) and those before 2008 as contemporary (n = 14). We acknowledge that using a three-year span of time requires us to infer syntopy; however, we included only five sites that required this inference. These sites were included due to our personal knowledge of the long-term presence of both species in these watersheds but where few reported observations exist. We also acknowledge personal knowledge of extant available and appropriate habitat at these sites. All other sites included contemporaneous occurrence of both species from 1 to 12 years (mean = 1.93 yrs).

Our own field research produced six locations where we directly identified both species co-occurring at a site between 2 and 12 years (**Table 1**). Foothill yellow-legged frogs were detected in one stock pond (i.e., Turtle Pond, Mitsui Ranch, Sonoma County) and were syntopic with a large population of California red-legged frogs using the pond (Alvarez and Wilcox 2021a). Foothill yellow-legged frogs were also observed in several creeks that are typically intermittent or ephemeral. In other areas, such as Copeland Creek in Sonoma County, both species were regularly present at the same site in the same pools, at the same time period (Erway 2022; Wilcox, unpublished data). This frequency of co-occurrence is similar in Napa County at Wragg Creek where a large pool (20 m x 20 m x 2 m deep) was occupied by both species when the pool was full. When the same pool dried, presumably both species occupied similar upland habitat for refuge sites adjacent to the creek (pers. obs.). We visited a site in Baja California at an elevation of 2,075 m where Loomis (1965) reported collecting two foothill-yellow legged frogs. At the time of our visit (58 years later), the habitat appeared as described by Loomis but only California red-legged frogs were present and were abundant. Further, Welsh (1988) also reported no indication of foothill yellow-legged frogs in this area and considered them marginal species in this region. Our visits suggest that habitat is present, but we were not able to conduct night surveys which would have elucidated presence or lack of presence more clearly (Alvarez and Wilcox, in press).

We found six contemporary observations that were reported through the CNDDB that were syntopic (**Table 1**). An additional 16 sites were considered historic (reported >15 years ago), with habitat that appears to remain intact and therefore may potentially support both species currently. We also had eight additional personal observations of sites that were contemporary, which totals 14 separate recent sites from five counties where the species can be found at the same site, in the same habitat, and at the same time (**Table 1**). Although our contemporary data are lacking observations in the Sierra Nevada, Barry and Fellers (2013) suggested that California red-legged frogs and foothill yellow-legged frogs were possibly syntopic in the Sierra Nevada. However, little work has been done in the Sierra Nevada largely due to immense private property ownership where the two species' ranges overlap.

In our compiled data, it was uncommon to detect both species in the same water body but enough to prompt us to investigate whether this is a common occurrence over a larger geographic area (i.e., multiple counties or hydrologic units). We suggest that where these species are sympatric, they are occasionally syntopic. Since the California red-legged frog is an anuran-eating species (Hayes and Tennant 1985; Alvarez 2013), and foothill yellow-legged frogs are a significantly smaller species, especially in the post-metamorphic stage, this may pose a threat to the foothill yellow-legged frogs

(McGinnis and Stebbins 2018; Alvarez et al. 2022). However, in a recent study where the two species are syntopic in Copeland Creek, Sonoma County, Erway (2022) reported that niche separation may be driven by frog size. Smaller frogs of both species remain close the wetted edge of the creek pools, while large adult California red-legged frogs are able to bask and forage further from the wetted edge because their surface-to-volume ratio allows them to thermoregulate more efficiently, therefore maintaining better hydration away from the water.

Although detected in stock ponds (Wilcox and Alvarez 2019; Alvarez and Wilcox 2021a), we recognize that these species did not naturally occur in ponds to the extent that we see them today because natural ponds were uncommon prior to European settlement (Schoenherr 1976). Thus, California red-legged frogs certainly evolved in lotic systems, and these were likely ephemeral systems that had intermittent pools that were suitable for oviposition and larval development. Since both species evolved in and were adapted for lotic habitats, it is reasonable to think both species historically overlapped in these systems.

Management concerns such as managed hydroperiod, extent of emergent or riparian vegetation, presence of potential predators, or breeding frequency may also vary between the two species. Nonetheless, our observations suggest that aquatic breeding habitat that is preserved, restored, or created within the range of both species—although frequently developed and managed for only one species—may well be used by both. This might include re-vegetating the margins or recontouring slopes adjacent to lotic habitats for restoration that may ultimately alter suitability for one or more species (Alvarez and Wilcox 2021b). The habitat requirements of both species should be carefully considered when developing or managing aquatic breeding sites, particularly low gradient streams within the overlapping range of these amphibians.

Our data indicate a pattern of syntopy among these two declining ranids within various aquatic habitats and over a large area—five counties currently with five additional counties historically—within their overlapping geographic ranges. Additional studies may further validate and expand the pattern we observed here.

Acknowledgments

We gratefully acknowledge The Sonoma Mountain Ranch Preservation Foundation, The Land Trust of Napa County, The East Bay Regional Park District, and California State Parks, for access to sites they manage. Joe Drennan of Kleinfelder generously shared information on the lifespan of foothill yellow-legged frogs. The Sonoma Mountain Ranch Preservation Foundation, The Wildlife Project, and Becky Rozumowicz-Kodsunit at Area West Environmental, Inc, supported this project in numerous ways and added to its success. We are also grateful to two anonymous reviewers who thoroughly reviewed an early draft of the manuscript and added greatly to its readability and clarity.

Literature Cited

- Allaback, M. L., D. M. Laabs, D. S. Keegan, and J. D. Harwayne. 2010. *Rana draytonii* (California red-legged frog). Dispersal. Herpetological Review 41:204.

- Alvarez, J. A. 2013. *Rana draytonii* (California red-legged frog). Cannibalism. Herpetology Review 44:126–127
- Alvarez, J. A., M. A. Shea, J. T. Wilcox, M. L. Allaback, S. M. Foster, G. E. Padgett-Flohr, and J. L. Haire. 2013. Sympatry in the California tiger salamander and California red-legged frog breeding habitat within their overlapping range. California Fish and Game 99:42–48.
- Alvarez, J. A. , L. L. Surber, E. A. Gallanty, and J. T. Wilcox. 2022.Small size of *Rana boylii* (Anura: Ranidae) at metamorphosis in a seasonal stream may confound survey methods. Herpetology Notes 15:395–397.
- Alvarez, J. A., and J. T. Wilcox. 2021a. Observations of atypical habitat use by foothill yellow legged frogs (*Rana boylii*) in the coast range of California. Western North American Naturalist 81:293–299.
- Alvarez, J. A., and J. T. Wilcox. 2021b. Observations of nocturnal upland habitat use by the *Rana draytonii* (California red-legged frog), and implications for restoration and other activities. Ecological Restoration 39:155–157.
- Alvarez, J. A., and J. T. Wilcox. In press. An effective method for long-term control of American bullfrogs syntopic with special-status native amphibians in California. Human-Wildlife Interactions.
- Barry, S. J., and G. M. Fellers. 2013. History and status of the California red-legged frog *(Rana draytonii*) in the Sierra Nevada, California, USA. Herpetological Conservation and Biology 8:456–502.
- Bondi, C. A., S. M. Yarnell, and A. J. Lind. 2013. Transferability of habitat suitability criteria for a stream breeding frog (*Rana boylii*) in the Sierra Nevada, California. Herpetological Conservation and Biology 8:88–103.
- Bourque, R. M. 2008. Spatial ecology of an inland population of the foothill yellow-legged frog (*Rana boylii*) in Tehama County, California. Thesis, Humboldt State University, Arcata, CA, USA.
- Bulger., N. J. Scott, Jr., R. B. Seymour, R. B. 2003. Terrestrial activity and conservation of adult California red-legged frogs (*Rana aurora draytonii*) in coastal forests and grasslands. Biological Conservation 110:85–95.
- California Fish and Game Commission (CFGC). 2020. Notice of findings for foothill yellow-legged frog (*Rana boylii*) – March 10, 2020. California Fish and Game Commission, Sacramento, CA, USA. Available from: **https://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=177905#:~:text=The%20Commission%20determined%20that%20the,management%20efforts%20required%20by%20CESA**.
- Cook, D. G., and M. R. Jennings. 2007. Microhabitat use of the California red-legged frog and introduced bullfrog in a seasonal marsh. Herpetologica 63:430–440.
- Erway, A. L. 2022. Comparative microhabitat use of two California native ranids, California red-legged frog (*Rana draytonii*) and foothill yellow-legged frog (*Rana boylii*), in a riparian woodland. Thesis, Sonoma State University, Rohnert Park, CA, USA.
- Feaver, P. E. 1971. Breeding pool selection and larval mortality of three California amphibians: *Ambystoma tigrinum californiense* Gray; *Hyla regilla* Baird and Girard; and *Scaphiopus hammondi hammondi* Girard. Thesis, Fresno State College, Fresno, CA, USA.
- Flaxington, W. C. 2021. Amphibians and Reptiles of California: Field Observations, Distribution, and Natural History. Fieldnotes Press, Anaheim, CA, USA.
- Hayes, M. P., and M. R. Jennings. 1988. Habitat correlates of distribution of the California red-legged frog (*Rana aurora draytonii*) and the foothill yellow-legged frog (*Rana boylii*): implications for management. Pages 144–158 in R.C. Szaro, K. E. Severson, and D. R. Patton, editors. Proceedings of the Symposium on the Management of Amphibians, Reptiles, and Small Mammals in North America. General Technical Report RM-166, USDA Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO, USA.
- Hayes. M. P., and M. R. Tennant. 1985. Diet and feeding behavior of the California red-legged frog *Rana aurora draytonii* (Ranidae). Southwestern Naturalist 30:601–605.

- Jennings, M. R., and M. P. Hayes. 1994. Amphibian and Reptile Species of Special Concern in California. California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova, CA, USA.
- Lannoo, M., editor. 2005. Amphibian Declines: The Conservation Status of United States Species. University of California Press, Berkeley, CA, USA.
- Loomis, R. B. 1965. The yellow-legged frog, (*Rana boylii*) from the Sierra San Pedro Martir, Baja, California Norte, México. Herpetologica 21(1):78–80.
- McGinnis, S. M., and R. C. Stebbins. 2018. Field Guide to the Amphibians and Reptiles of California. 4th edition. University of California Press, Berkeley, CA, USA.
- Peralta-García, A., J. A. Alvarez, J. H. Valdez-Villavicencio, and B. D. Hollingsworth. 2022. Longevity in wild California red-legged frogs (*Rana draytonii*) from Baja California, México. Bulletin of the Southern California Academy of Sciences 121:1–4.
- Petranka, J. W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, D.C., USA.
- Rathbun, G. B. 1998. *Rana aurora draytonii* (California red-legged frog). Egg predation. Herpetological Review 29:165.
- Schoenherr, A. A. 2007. A Natural History of California. 2nd edition. University of California Press, Berkeley, CA, USA.
- Stebbins, R. C. 2003. A Field Guide to Western Reptiles and Amphibians. 3rd edition. Houghton Mifflin Company, Boston, MA, USA.
- Stebbins, R. C., and S. M. McGinnis. 2012. A Field Guide to Amphibians and Reptiles of California. University of California Press, Berkeley, CA, USA.
- Storer, T. I. 1925. A synopsis of the amphibia of California. University of California Publications in Zoology 27:1–342.
- Thomson, R. C., A. N. Wright, and H. B. Shaffer. 2016. California Amphibian and Reptile Species of Special Concern. University of California Press, Berkeley, CA, USA.
- U.S. Fish and Wildlife Service (USFWS). 2022. Endangered and Threatened Wildlife and Plants; foothill yellow-legged frog; threatened status with Section 4(d) Rule for two distinct population segments and endangered status for two distinct population segments. 202286 FR:73914–73945.
- Welsh, H. H., Jr. 1988. An ecogeographic analysis of the herpetofauna of the Sierra San Pedro Mártir region, Baja, California, with a contribution to the biogeography of the Baja California herpetofauna. Proceedings of the California Academy of Sciences 46:1–72.
- Wilcox, J. T., and J. A. Alvarez. 2019. Wrestling for real estate: male-male interactions in breeding foothill yellow-legged frogs (*Rana boylii*). Western Wildlife 6:14–17.
- Zweifel, R. G. 1955. Ecology, distribution , and systematics of frogs of the *Rana boylei* group. University of California Publications in Zoology 54:207–292.